








How twisty is that orbit?

Anand Deopurkar

2024

NZMS AUSTMS AMS






Orbits of group actions


G G V

\overline{Gv} \leftarrow v




$$G = \mathbb{G}_m^2$$


$$V = \mathbb{A}^3$$




$(1,1), (1,2), (2,1)$




$(1,1), (1,2), (3,1)$




$(3,1), (2,2), (1,3)$



$(3,2), (4,4), (1,3)$



$(5,1), (1,2), (3,1)$




$(5,1), (4,2), (3,5)$

$$G = GL(2)$$

$$V = \text{Sym}^4(2)$$

$$v = XY(X-Y)(X-3Y)$$




(3d slice)

$$G = GL(2)$$

$$V = \text{Sym}^4(2)$$

$$v = XY(X-Y)(X+Y)$$




(3d slice)

Orbits of group actions

$$G \curvearrowright V$$

$$X = \overline{Gv} \quad \leftarrow \quad v$$



1) $\dim X = ?$


2) $\deg X = ?$

3) $\text{Sing } X = ?$


4) $\partial X = (\overline{Gv} - Gv) = ?$

Geometric ramifications

Ex: $G = GL(2)$
 $V = \text{Sym}^4(2)$

$v =$ 
 $Gv = \left\{ \begin{array}{l} \text{4-tuples with the} \\ \text{same cross-ratio} \end{array} \right\}$

$G = GL(3)$
 $V = \text{Sym}^3(3)$

$v =$ 
 $Gv = \left\{ \begin{array}{l} \text{Elliptic curves with the} \\ \text{same } j\text{-invariant} \end{array} \right\}$

Other groups,
other representations

\Rightarrow Other interpretations.

What is known?

- $GL(2)$, $\text{Sym}^n(2)$ Enriques-Fano 1890s
- $GL(3)$, $\text{Sym}^n(3)$ Aluffi-Faber 1990s
- $GL(4)$, $\text{Sym}^3(4)$ D-Patel-Tseng 2021
 - ↳ "27 questions about cubic surfaces" by Ranestad-Sturmfels
- G_m^n , any rep D. 2024
- $GL(2)$, any rep D. 2024
- Reps for quiver moduli Berget, Buch, Fink, Feher (2000s)
Fulton, Rimanyi, Weber

How?




Orbit
degrees



Equivariant
fundamental classes

Equivariant Fundamental Class

$$X = \overline{Gv} \leftarrow \begin{matrix} G & G & V \\ & \leftarrow & v \end{matrix}$$



$$\deg X = [X] \in H^*(\mathbb{P}V) \leftarrow \mathbb{Z}$$

$$Efc(X) = [X] \in H_G^*(V) \leftarrow \text{Polynomial ring}$$

References

1) A universal formula for counting cubic surfaces

DEOPURKAR, PATEL, TSENG,
To appear in Algebraic Geometry

2) Equivariant classes of orbits in $GL(2)$ representations

DEOPURKAR, Pre-print (arxiv)

3) Orbits of linear series on the projective line

DEOPURKAR, PATEL

International Mathematical Research Notices (2024)

From: Anand Deopurkar <anand.deopurkar@anu.edu.au>

Subject: **Exciting! Re: Update on cubic surfaces**

To: Anand Patel <anandppatel@gmail.com>, Dennis Tseng <ehhheehee@gmail.com>

Date: Wed, 02 Jun 2021 11:38:30 +1000 (3 years, 26 weeks, 2 days ago)

Hi Anand and Dennis,

I followed yesterday's line of thinking to its conclusion. From the GIT of (X,H) , we can write down the following cubic surfaces with G_m actions that are provably not in the closure of a general smooth cubic.

1. $x_0x_2x_3 = x_1^3$
2. $x_3(x_0x_2 - x_1^2) = x_0x_1^2$
3. $x_3(x_0x_2 - x_1^2) = x_0^2x_1$
4. $x_0^2x_3 = x_1x_2(x_1 + x_2)$
5. $x_0x_2^2 = x_1^3 + x_0x_1x_3$

Each one gives a relation (the last one gives the same relation as the first). Together with the relation written on Overleaf for the family B_1 , namely $16a_1^4 + 4a_1^2a_2 + a_2^2 = 4320$, we can write down the following (integral) relation:

$$1080 * a_1^2a_2 - 1080 * a_1a_3 + 9720 a_4 = \text{Cubic surface count.}$$

Amazingly, this agrees with the 96120 count for the family given by a general web

THANK YOU!