

# Degree

$F$  a field,  $F \subset K$   $K$  also a field.

$\deg_F K = \deg(K/F) := \dim$  of  $K$  as an  $F$  v. space.

If finite, say that  $K$  is a finite ext<sup>n</sup> of  $F$ .

Ex.  $\alpha$  is an algebraic complex number  
then  $\mathbb{Q}[\alpha]$  is a finite ext<sup>n</sup> of  $\mathbb{Q}$ .  
& its deg = deg of min poly of  $\alpha$ .

Notation:

$\mathbb{Q}(\alpha) =$  Smallest subfield of  $\mathbb{C}$  containing  $\mathbb{Q}$  &  $\alpha$

$\alpha$  algebraic  $\Rightarrow \mathbb{Q}(\alpha) = \mathbb{Q}[\alpha]$

$\alpha$  transc.

$\mathbb{Q}(\alpha) = \left\{ \frac{p(\alpha)}{q(\alpha)} \mid p, q \in \mathbb{Q}[x] \right\}$

## Multiplicative property:

$$\underbrace{F \subset K} \subset \underbrace{K \subset L}$$

fields.

Thm: If  $K$  finite over  $F$  &  $L$  finite over  $K$   
then  $L$  finite over  $F$  &

$$\deg_F L = \deg_K L \cdot \deg_F K$$

Pf idea: Say  $a_1, \dots, a_m$  is a basis of  $L$  over  $K$ .  
Say  $b_1, \dots, b_n$  is a basis of  $K$  over  $F$ .

Then  $\{b_i \cdot a_j, i=1, \dots, m, j=1, \dots, m\}$  is a  
basis of  $L$  over  $F$ .

Consequences:  $\mathbb{Q} \subset \mathbb{C}$

$\alpha, \beta \in \mathbb{C}$  algebraic over  $\mathbb{Q}$ .

Then  $\alpha + \beta$ ,  $\alpha\beta$  &  $\alpha/\beta$  are all alg. /  $\mathbb{Q}$ .

Pf: Consider

$$\underbrace{\mathbb{Q} \subset \mathbb{Q}[\alpha]}_{\text{finite}} \subset \underbrace{\mathbb{Q}[\alpha, \beta]}_{\text{finite}} \ni \begin{array}{l} \alpha + \beta \\ \alpha\beta \\ \alpha/\beta \\ \alpha - \beta \end{array}$$

(  $F \subset K$  finite &  $\gamma \in K$  has to be alg /  $F$ .  
 $F \subset F[\gamma] \subset K$  so  $F[\gamma]$  must be fin dim /  $F$ . )

If  $F \subset K$  has degree  $d$ .

Then every  $\alpha \in K$  has degree dividing  $d$ .

$$\underbrace{F \subset F[\alpha] \subset K.}$$

Ex:  $\sqrt[3]{2} \notin \mathbb{Q}[\sqrt{2}, \sqrt{3}, \sqrt{5}, \sqrt{17}]$

$\downarrow$

$X^3 - 2$  irred

$\Downarrow$

deg 3.

$\mathbb{Q}[\sqrt{2}, \sqrt{3}, \sqrt{5}, \sqrt{17}]$  } 2 or 1

$\cup$

$\mathbb{Q}[\sqrt{2}, \sqrt{3}, \sqrt{5}]$  } 2 or 1

$\cup$

$\mathbb{Q}[\sqrt{2}, \sqrt{3}]$  } 2 or 1

$\cup$

$\mathbb{Q}[\sqrt{2}]$  } 2 or 1

$\cup$

$\mathbb{Q}$  } 2 or 1


} 2<sup>mm</sup>

(\*)  $\implies$  will prove later

# Constructing using ruler & compass.



Ruler ← draws a line through 2 pts.



centre

Compass ↑

Constructed

Draws a circle with given center & through a given pt.

Game: Start with some pts. ←

- ① Draw a line through 2 constructed pts.
- ② Circle centered at a cons. pt through a cons. pt.
- ③ Intersection pts are constructed.


Problem : Given

$p$


$q$ .

Task : ~~Find~~ midpoint.

Construct.




Problem : Given



Want to construct a pt  $s$  so that  
 $\angle qpr$  is bisected by  $ps$ .

Problem: Trisect a segment.

Given



Want  $r, s$  that trisect it.

Can be done.

Problem: Trisect an angle.



cannot be  
done!